

The Lipid Solubility of Most Drugs Play Important Role of Its Pharmacological Action and Duration of Action

Rezk R. Ayyad¹, Ahmed M. Nejm¹, Yasser Abdel Allem Hassan² & Ahmed R. Ayyad³

¹ Pharmaceutical Medicinal Chemistry Department, Faculty of Pharmacy, ALAZHAR University, Cairo, Egypt

² Pharmaceutics and Pharmaceutical Technology Department, Faculty of Pharmacy, Delta University for Science and Technology, Gamasa, Addaqahlya, Egypt

³ Faculty of Medicine, Asfendiyarov, Kazakh, National Medical University (KazNMU)

Correspondence: Rezk R. Ayyad, Pharmaceutical Medicinal Chemistry Department, Faculty of Pharmacy, ALAZHAR University, Cairo, Egypt.

doi:10.56397/JPEPS.2023.12.01

Abstract

The fact of human body and animal body formed of organs which formed of tissue which its constituents are the cells.

The cell enveloped by cell membrane which contain more its content of lipid compounds to protect the cell. Hence when the chemical compounds (drugs) across to the cytoplasm of the cell must have a lipophilic character. Also there is a fact the oily substances not mix with the watery substance, so must be take a cream form (water in oil or oil in water emulsion), so the drugs must be soluble in lipid to across the cell to obtain the desirable effect.

The partition co-efficient is a significant expression of lipid solubility.

Many of drugs convert to prodrug this prodrug has highly lipid solubility and has long duration of action e.g., testosterone cypionate, rolitetracycline, penicillin procaine, benzylpenicillin G, nandrolone decanoate....etc., on the other hand the water soluble compounds (hydrophilic drugs) e.g., vitamin B complexes which exert its effects and excreted from the kidney quickly this indicate to no long duration of vitamin B complex except the cyanocobalamin (vit.B12) which incorporated to sesame oil to deposit in the body to treatment the inflammation of nerves of diabetic patients which suffering from neuropathy.

Keywords: lipid solubility, hydrophilic drugs, partition co-efficient, prodrugs, duration of action, pharmacological action, blood brain barrier (BBB)

1. Introduction

In the first when the chemist design new drug, or the physician want to describe some drugs in

some diseases must be taken in his mind the lipid solubility of the drug to sure reach the drug to target site and may be prolonged its action. The partition co-efficient is significant for lipid solubility in vitro where represented by the solubility of the drug in n-octane (represent the lipid phase), the water which represent the aqueous phase, if the chemical compound (drug) soluble in octane phase will be soluble in lipid and if soluble in water will be water soluble. The partition co-efficient is the ratio of drug soluble in n-octane to water. Many of drugs may be react with organic acids (acetic, propionic, butyric, pentanoic, hexanoic, heptanoic, octanoic, nonanoic, decanoic, undecanoic, dodecanoic) or fatty acids (saturated non-essential, non-saturated essential), to make esters, these esters don't water soluble (many of esters mainly water insoluble), so these esters lipid soluble not only this fat but the drugs become prodrugs. The lipid solubility of drug esters makes the drugs reach to the target sites due to lipid solubility across the cell membrane of (cell, tissue or organ), furthermore the drug esters easily hydrolyzed by esterase enzyme in the body. Hence the pharmacists, physicians, and the pharmaceutical companies must take in consideration the lipid solubility in your works to reach the maximum activity of the drugs.

2. Chemistry and Discussion

The drugs which are water soluble not stay in the body for long time e.g., water soluble vitamins where it excreted from the kidney. Vit.B1 (Thiamine).



Thiamine is important to growth and vital functions and present in many foods as (dicots) and is important for metabolism of carbohydrates, lipids and protein, decrease of the thiamine in the body will cause dysfunction on the body and lead to Beriberi disease, so we must take in consideration the solubility of thiamine and its excretion quickly from the kidney.

The Riboflavin is vit.B2 which is essential for the

functions of the body.



Riboflavin plays many important roles in vital processes in the body, the very important role is in carbohydrates (glucose) metabolism which enter the structure of FAD which is Co-enzyme to enzymes which act on glucose, also water soluble and excreted from the kidney.

Nicotinamide is vit.B3 one of the vit.B complex group which make vital role in skin health and its important role enter the structure NAD and NADP which are important as co-enzymes to the enzyme which act on glucose, so when decreased in the body due to its water solubility and excreted from kidney may be increase the glucose level in the body.



Pyridoxine is a vit.B6 and is also member of vit.B complex which is important for keep on the health nerves especially in diabetic patients which after time of chronicity wine neuropathy, also used in pregnant woman emesis (also has water solubility and excreted from the body via excretion routes).



Vit.B12 (cyanocobalamin) also one of the vit.B complex which soluble in water and decrease in the body, so it must be taken more times for diabetic and pre diabetic patients because it keep and treat the inflammation of nerves caused by high level of glucose in the blood. As we mentioned above the cyanocobalamin in sesame oil to prolonged in the body.



Vit.C (Ascorbic acid) is a water soluble vitamin plays an essential role in the body to maintain a healthy life as it act as powerful antioxidant, raised the immune level so enhance immunity against infection and help in relief the symptoms of common cold and other viral infections, also it enhance absorption of Ca so maintain a healthy bone, also ascorbic acid (vit.C) water soluble and excreted from the body and the body need take repeated administration to keep its function in body, all the mentioned vitamins are essential for human, but its water solubility make them make them around disadvantage.



Also Pantothenic acid (vit.B5), Biotin (vit.B7), Folic acid (vit.B9) and Amygdaline (vit.B17 which use as anticancer) these compounds are water soluble has low concentration in the body.



N.B. The vitamins which soluble in water must be alarm to physicians of diabetes and must calculate the actual need of vitamin B complex which are very important in diabetic treatment.

3. Conclusion

The lipid solubility must be taken in consideration in many drugs when we want design a new drug or the physician want to describe a drug to patient. Absolutely not all drugs are highly lipid solubility due to the accumulation property e.g., Digoxin not merely, some compounds if stored in the body may be harm to it (toxicity of digoxin) or cause disease (edema of mineralocorticoids hormone), in spite we attention to the importance of lipid solubility of the drugs to sure reach it to the target sites or to give prolonged action.



References

- A Ibrahim, HM Sakr, RR Ayyad, MM Khalifa. (2022). Design, Synthesis, In-Vivo Anti-Diabetic Activity, In-Vitro α -Glucosidase Inhibitory Activity and Molecular Docking Studies of Some Quinazolinone Derivatives. *ChemistrySelect*, 7(14), e202104590.
- AA El-Helby, MK Ibrahim, AA Abdel-Rahman, RRA Ayyad, MA Menshawy, ... (2009). Synthesis, molecular modeling and anticonvulsant activity of benzoxazole derivatives. *Al-Azhar J Pharm Sci*, 40, 252-70.
- AA Elhelby, RR Ayyad, MF Zayed. (2011). Synthesis and biological evaluation of some novel quinoxaline derivatives as anticonvulsant agents. *Arzneimittelforschung*, *61*(07), 379-381.
- AAM Abdel-Aziz, AS El-Azab, AM Alanazi, YA Asiri, IA Al-Suwaidan, ... (2016). Synthesis and potential antitumor activity of 7-(4-substituted

piperazin-1-yl)-4-oxoquinolines based on ciprofloxacin and norfloxacin scaffolds: *in silico* studies. *Journal of Enzyme Inhibition and Medicinal Chemistry*, *31*(5), 796-809.

- AGA El-Helby, H Sakr, RR Ayyad, HA Mahdy, MM Khalifa, A Belal, ... (2020). Design, synthesis, molecular modeling, in vivo studies and anticancer activity evaluation of new phthalazine derivatives as potential DNA intercalators and topoisomerase II... *Bioorganic chemistry*, 103, 104233.
- AGA El-Helby, H Sakr, RRA Ayyad, K El-Adl, MM Ali, F Khedr. (2018). Design, synthesis, in vitro anti-cancer activity, ADMET profile and molecular docking of novel triazolo [3, 4-a] phthalazine derivatives targeting VEGFR-2 enzyme. *Anti-Cancer Agents in Medicinal Chemistry (Formerly Current Medicinal Chemistry-Anti-Cancer Agents)*.
- AGA El-Helby, H Sakr, RRA Ayyad, K El-Adl, MM Ali, F Khedr. (2018). Anti-Cancer Agents Med. *Chem*, *18*(8), 1184.
- AGA El-Helby, RR Ayyad, HM Sakr, AS Abdelrahim, K El-Adl, FS Sherbiny, ... (2017). Design, synthesis, molecular modeling and biological evaluation of novel 2, 3-dihydrophthalazine-1, 4-dione derivatives as potential anticonvulsant agents. *Journal of Molecular Structure*, 1130, 333-351.
- AGA El-Helby, RRA Ayyad, H Sakr, K El-Adl, MM Ali, F Khedr. (2017). design, synthesis, molecular docking, and anticancer activity of phthalazine derivatives as VEGFR-2 inhibitors. *Archiv der Pharmazie*, 350(12), 1700240.
- AGA El-Helby, RRA Ayyad, K El-Adl, A Elwan. (2017). Quinoxalin-2(1H)-one derived AMPA-receptor antagonists: Design, synthesis, molecular docking and anticonvulsant activity. *Medicinal Chemistry Research*, 26, 2967-2984.
- AGA El-Helby, RRA Ayyad, K El-Adl, H Elkady. (2019). Phthalazine-1, 4-dione derivatives as non-competitive AMPA receptor antagonists: design, synthesis, anticonvulsant evaluation, ADMET profile and molecular docking. *Molecular diversity*, 23, 283-298.
- AGA El-Helby, RRA Ayyad, K El-Adl, H Sakr, AA Abd-Elrahman, IH Eissa, ... (2016). Design, molecular docking and synthesis of some novel 4-acetyl-1-substituted-3,4-dihydroquinoxali n-2(1*H*)-one derivatives for anticonvulsant evaluation as AMPA... *Medicinal Chemistry Research*, 25, 3030-3046.

- AGA El-Helby, RRA Ayyad, MF Zayed, HS Abulkhair, H Elkady, K El-Adl. (2019). Design, synthesis, in silico ADMET profile and GABA-A docking of novel phthalazines as potent anticonvulsants. *Archiv Der Pharmazie*, 352(5), 1800387.
- AM Alaa, AS El-Azab, AM Alanazi, YA Asiri, IA Al-Suwaidan, AR Maarouf ... (2015). Synthesis and potential antitumor activity of 7-(4-substituted piperazin-1-yl)-4-oxoquinolines based on ciprofloxacin and norfloxacin scaffolds: in silico studies. *Journal of Enzyme Inhibition and Medicinal Chemistry*.
- AM Alaa, AS El-Azab, LA Abou-Zeid, KEH ElTahir, NI Abdel-Aziz, ... (2016). Synthesis, anti-inflammatory, analgesic and COX-1/2 inhibition activities of anilides based on 5, 5-diphenylimidazolidine-2, 4-dione scaffold: molecular docking studies. *European journal* of medicinal chemistry, 115, 121-131.
- AM Alanazi, AAM Abdel-Aziz, TZ Shawer, RR Ayyad, AM Al-Obaid, ... (2016). Synthesis, antitumor and antimicrobial activity of some new 6-methyl-3-phenyl-4(3*H*)-quinazolinone analogues: *in silico* studies. *Journal of Enzyme Inhibition and Medicinal Chemistry*, 31(5), 721-735.
- AS El-Azab, AM Alaa, RR Ayyad, M Ceruso, CT Supuran. (2016). Inhibition of carbonic anhydrase isoforms I, II, IV, VII and XII with carboxylates and sulfonamides incorporating phthalimide/phthalic anhydride scaffolds. *Bioorganic & medicinal chemistry*, 24(1), 20-25.
- E Nassar, YA El-Badry, AMM Eltoukhy, RR Ayyad. (2016). Synthesis and Antiproliferative Activity of 1-(4-(1H-Indol-3-Yl)-6-(4-Methoxyphenyl) Pyrimidin-2-yl) Hydrazine and Its Pyrazolo Pyrimidine Derivatives. *Med chem (Los Angeles)* 6, 224-233.
- H M Sakr, R R Ayyad, K Mahmoud, A M Mansour, G Ahmed. (2021). Design, Synthesis of Analgesics and Anticancer of Some New Derivatives of Benzimidazole. *International Journal of Organic Chemistry*, 11(03), 144-169.
- H Mahdy, M Shaat, R Ayyad. (2022). RECENT ADVANCES IN DRUGS TARGETING PROTEIN KINASES FOR CANCER

THERAPY. *Al-Azhar Journal of Pharmaceutical Sciences*, 66(2), 56-86.

- H Sakr, RR Ayyad, AA El-Helby, MM Khalifa, HA Mahdy. (2021). Discovery of novel triazolophthalazine derivatives as DNA intercalators and topoisomerase II inhibitors. *Archiv der Pharmazie*, 354(6), 2000456.
- IA Al-Suwaidan, AAM Abdel-Aziz, TZ Shawer, RR Ayyad, AM Alanazi, ... (2016). Synthesis, antitumor activity and molecular docking study of some novel 3-benzyl-4 (3H) quinazolinone analogues. *Journal of enzyme inhibition and medicinal chemistry*, 31(1), 78-89.
- IA Osman, RR Ayyad, HA Mahdy. (2022). New pyrimidine-5-carbonitrile derivatives as EGFR inhibitors with anticancer and apoptotic activities: design, molecular modeling and synthesis. *New Journal of Chemistry*, 46(24), 11812-11827.
- IH Eissa, AM Metwaly, A Belal, ABM Mehany, RR Ayyad, K El-Adl, ... (2019). Discovery and antiproliferative evaluation of new quinoxalines as potential DNA intercalators and topoisomerase II inhibitors. *Archiv der Pharmazie*, 352(11), 1900123.
- K El-Adl, AGA El-Helby, H Sakr, RR Ayyad, HA Mahdy, M Nasser, ... (2021). Design, synthesis, molecular docking, anticancer evaluations, and in silico pharmacokinetic studies of novel 5-[(4-chloro/2, 4-dichloro) benzylidene] thiazolidine-2, 4-dione... *Archiv der Pharmazie*, 354(2), 2000279.
- K El-Adl, AGA El-Helby, RR Ayyad, HA Mahdy, MM Khalifa, HA Elnagar, ... (2021). Design, synthesis, and anti-proliferative evaluation of new quinazolin-4 (3H)-ones as potential VEGFR-2 inhibitors. *Bioorganic & Medicinal Chemistry, 29*, 115872.
- M Al Ward, AE Abdallah, M Zayed, R Ayyad, M El-Zahabi. (2023). New immunomodulatory anticancer quinazolinone based thalidomide analogs: Design, synthesis and biological evaluation.
- M Salem, R Ayyad, H Sakr. (2022). Design and Synthesis of Some New Oxadiazole Derivatives as Anticancer Agents. *International Journal of Organic Chemistry*, 12(02), 64-74.
- MF Zayed, RR Ayyad. (2012). Some novel anticonvulsant agents derived from phthalazinedione. *Arzneimittelforschung*,

62(11), 532-536.

- MK Ibrahim, AA Abd-Elrahman, RRA Ayyad, K El-Adl, AM Mansour, ... (2013). Design and synthesis of some novel 2-(3-methyl-2-oxoquinoxalin-1 (2H)-yl)-N-(4-(substituted) phenyl) acetamide derivatives biological for evaluation as anticonvulsant agents. Bulletin of Faculty of Pharmacy, Cairo University, 51(1), 101-111.
- MK Ibrahim, AEA El-Helby, AH Ghiaty, AH Biomy, AA Abd-El Rahman. (2009). Modeling, Synthesis and Antihyperglycemic Activity of Novel Quinazolinones Containing Sulfonylurea. J. Biol. Pham. Sci., 7(1).
- MM Khalifa, HM Sakr, A Ibrahim, AM Mansour, RR Ayyad. (2022). Design and synthesis of new benzylidene-quinazolinone hybrids as potential anti-diabetic agents: In vitro α -glucosidase inhibition, and docking studies. *Journal of Molecular Structure*, 1250, 131768.
- R Ayyad, H Sakr, A Gaafer. (2022). Design and Synthesis of New Compounds Derived from Phenyl Hydrazine and Different Aldehydes as Anticancer Agents. *International Journal of Organic Chemistry*, 12(1), 28-39.
- R Ayyad. (2012). Synthesis and Biological Evaluation of Novel Iodophthalazinedione Derivatives as Anticonvulsant Agents. *Al-Azhar Journal of Pharmaceutical Sciences*, 45(1), 1-13.
- R Ayyad. (2014). Synthesis and Anticonvulsant Activity of 6-Iodo Phthalazinedione Derivatives. *Al-Azhar Journal of Pharmaceutical Sciences*, 50(2), 43-54.
- RA Ayyad, HM Sakr, KM El-Gamal. (2016). Design, Synthesis, Computer Modeling and Analgesic Activity of Some New Disubstituted Quinazolin-4 (3H)-ones. *Med. Chem*, 6(5), 299-305.
- RR Ayyad, AM Nejm, AR Ayyad. (2023). The Activity of Some Antibiotics Depend on Stereochemistry of Them (Its Structure). *Journal of Progress in Engineering and Physical Science*, 2(2), 5-7.
- RR Ayyad, AM Nejm, AR Ayyad. (2023). The Isomers of Some Drugs One Effective and the Other Is Toxic or Ineffective. *Current Research in Medical Sciences*, 2(2), 58-62.

- RR Ayyad, AM Nejm, YH Abdelaleem, AR Ayyad. (2023). Hydrophobicity, Transport and Target Sites of Action Are Important for the Activity of Many Drugs. *Current Research in Medical Sciences*, 2(3), 15-19.
- RR Ayyad, HM Sakr, KM El-Gamal, IH Eissa, A HA, AS Tita, FF Sherbini, ... (2017). Anti-Inflammatory, Proton Pump Inhibitor and Synthesis of Some New Benzimidazole Derivatives. *Der Chemica Sinica*, 8(1), 184-97.
- RRA Ayyad, H Sakr, K El-Gamal. (2016). Synthesis, modeling and anticonvulsant activity of some phthalazinone derivatives. *American Journal of Organic Chemistry*, 6(1), 29-38.
- T Al-Warhi, AM El Kerdawy, N Aljaeed, OE Ismael, RR Ayyad, ... (2020). Synthesis, biological evaluation and in silico studies of certain oxindole–indole conjugates as anticancer CDK inhibitors. *Molecules*, 25(9), 2031.
- T Al-Warhi, H Almahli, RM Maklad, ZM Elsayed, MA El Hassab, ... (2023).
 1-Benzyl-5-bromo-3-hydrazonoindolin-2-on es as Novel Anticancer Agents: Synthesis, Biological Evaluation and Molecular Modeling Insights. *Molecules*, 28(7), 3203.
- WM Eldehna, MF Abo-Ashour, T Al-Warhi, ST Al-Rashood, A Alharbi, ... (2021). Development of 2-oxindolin-3-ylidene-indole-3-carbohydraz ide derivatives as novel apoptotic and anti-proliferative agents towards colorectal cancer cells. *Journal of Enzyme Inhibition and Medicinal Chemistry*, 36(1), 320-329.
- WM Eldehna, R Salem, ZM Elsayed, T Al-Warhi, HR Knany, RR Ayyad, ... (2021). Development of novel benzofuran-isatin conjugates as potential antiproliferative agents with apoptosis inducing mechanism in Colon cancer. *Journal of Enzyme Inhibition and Medicinal Chemistry*, 36(1), 1423-1434.