

Organophosphorus Residues in Fish in Rural Areas

Saad M. Saad¹, Fahim A. Shaltout¹, Amal A. A. Farag¹ & Hashim F. Mohammed²

¹ Department of Food Control, Faculty of Veterinary Medicine, Benha Univerisity, Egypt

² Animal Health Research Institute, Dokki Branch, Giza

Correspondence: Fahim A. Shaltout, Department of Food Control, Faculty of Veterinary Medicine, Benha Univerisity, Egypt.

doi:10.56397/JPEPS.2022.11.05

Abstract

A total 45 fish samples (Tilapia, Claris) were collected from EL-Bagoria, EL-Menofi drainage and Bahr Shibin for detection and determination of organophosphorus pesticides (diazinon, Malathion and chlorpyrifos). Organophosphorus pesticides could not be detected in Bahr Shibin.

No diazinon in Tilapia from EL-Bagoria but mean value of of Malathion 5.50 ± 2.0 , chlorpyrifos 5.0 ± 1.66 but from EL-Menofi drainage mean level of diazinon 3.33 ± 0.65 , Malathion 4.98 ± 1.22 and chlorpyrifos 3.0 ± 0.71 but in Claris samples, there were no diazinon, no Malathion, no chlorpyrifos from EL-Bagoria canal but from EL-Menofi drainage, mean value of diazinon 6.8 ± 1.42 , Malathion 4.64 ± 0.81 and chlorpyrifos 4.0 ± 0.96 .

Pesticides are one the main contaminant of water sources which are considered the natural environment of fish. on the other hand, fish could be contaminated by pesticides either directly by gills breathing or indirectly through contamination of feeding items, so it is very important to analyze fish samples to detect to what extent the rate of accumulation of pesticides residues in fish flesh and organs.

Keywords: Organophosphorus, diazinon, Malathion, chlorpyrifos

1. Introduction

Fish is low-fat high-quality protein. Fish is filled with omega-3. Fatty acids and vitamins such as D and B2 (riboflavin). Fish is rich in calcium and phosphorus and great source of iron, zinc, iodine, magnesium and potassium. The American Heart Association recommends eating fish at least two times per week as part of a healthy diet. Fish is packed with protein, vitamins and nutrients that can lower blood pressure and help to reduce the risk of a heart attack or stroke.

Fish are known to be highly nutritious and excellent sources of animal protein which are consumed by larger percentage of the world's population because of its availability and palatability (Shaltout, 2003; Hassan et al., 2014; Edris et al., 2017a). Fish and fish products are in the forefront of food safety and quality improvement because they are among the most internationally traded food commodities. Fish and fish products are one of the most important food stuffs as they are one of the cheapest sources of

animal protein. Fish are enriched with essential minerals, vitamins and unsaturated fatty acids (Shaltout & Hashim 2002; Hassan & Shaltout, 2004; Shaltout et al. 2015; Edris et al., 2017 b; Hassan et al., 2019). Today, the environmental pollution is considered one of the most serious problems in the world. The deleterious effect of the environmental pollution of pesticides is one of the principal research activities 1962. Contamination of food at animal origin with Organochlorine compounds and their metabolites has been reported in various countries (Neumann, 1988; Goldman et al., 1990).

Choudhary et al. (2014): reported that Organophosphorus pesticides causes burning/stinging of eyes, blurred vision, skin redness and itching, excessive sweating and shortness of breath, dry sore throat and burning of nose among spray farmers of Bhopal, Madhya Pradesh India, who sprayed pesticides by themselves and therefore were directly exposed to pesticides.

When people come into contact with large quantities of pesticide, this may cause acute poisoning or long-term health effects, including cancer and adverse effects on reproduction (FAO, WHO, 1987).

Fish and aquatic animals are exposed to pesticides in three primary ways, the first way was through dermal by direct absorption through the skin by swimming in pesticide contaminated waters, the second is breathing by direct uptake of pesticide through the gills during respiration and the third way is oral by drinking pesticide contaminated water or feeding on pesticide contaminated preys. Poisoning of human being occur by consumption of poisoned fish, which termed (secondary poisoning) Johnson and Finley (1980).

Priyadarshini et al. (2017): reported that

indiscriminate use and improper handling of synthetic pesticides in agriculture have resulted in serious problems such as asthma, wheeze and chronic bronchitis among the farmers.

2. Aim of Work

This study is made for detection and determination organophosphorus in Nile Tilapia and Claris samples from EL-Bagoria canal, EL-Menofi drainage and Bahr Shibin in Menofia Governorate.

3. Materials and Methods

45 fish samples of Tilapia and Claris from EL-Bagoria canal, EL-Menofi drainage and Bahr Shibin in Menofia Governorate for detection and determination of organophosphorus (Malathion, Diazinon and chlorpyrifos).

The collected samples were packed separately and transferred to laboratory for analysis.

Samples were extracted according to AOAC 1980. and Pesticide Analytical Manual 1978.

Fifty grams of samples were grinded with 100 gram of anhydrous sodium sulphate in presence of 150gram of 40-60petroleum ether for 2 minutes then extract was decanted through 500ml Buchnov funnel containing two wattman filter papers number 1/2.

The extract was poured through 40x25mm column of anhydrous sodium sulphate and eluent was collected in 500ml flask. Extraction and clean up by acetonitrile partitioning. Clean up by florisil column was firstly carried out by eluting the column with 200ml of 50 petroleum ether /diethyl ether (v/v). The elute was concentrated by rotator evaporator to dry film which was dissolved by 2ml n-hexane for HPLC determination.

4. Results

Table 1. Statistical analysis of organophosphorus residues (ppb, wet weight) in examined Tilapia samples

Pesticides	Positive samples in Bahr Shibin		Positive samples in El-Bagoria canal		Positive samples in El-Menofi drainage	
	No.	%	No.	%	No.	%
Diazinon	0	0%	0	0%	3	20%
Malathion	0	0%	2	13.3%	5	33.3%
chlorpyrifos	0	0%	3	20%	4	26.6%

Table 2. Mean residue levels of Organophosphorus pesticides (ppb, wet weight) in examined *Tilapia* samples from EL-Bagoria canal and EL-Menofi drainage

	Diazinon				Malathion				Chlorpyrifos			
	Min	Max	Mean	St. error	Min	Max	mean	St. error	Min	Max	mean	±St. error
El-Bagoria	0	0	0.00	0.00	3.5	7.5	5.50	2.0	2.75	8.25	5.0	1.66
EL-Menofi	2.25	4.5	3.33	0.65	2.25	9.15	4.98	1.22	1.5	4.75	3.0	0.71




Figure 1. Mean residue levels of Organophosphorus pesticides (ppb, wet weight) in examined *Tilapia* samples from EL-Bagoria canal and EL-Menofi drainage

Table 3. Statistical analysis of organophosphorus residues (ppb, wet weight) in examined Claris samples

Pesticides	Positive samples in Bahr Shubin		Positive samples in EL-Bagoria canal		Positive samples in EL-Menofi drainage	
	No.	%	No.	%	No.	%
Diazinon	0	0%	0	0%	6	40%
Malathion	0	0%	0	0%	9	60%
chlorpyrifos	0	0%	0	0%	5	33.3%

Table 4. Mean residue levels of Organophosphorus pesticides (ppb, wet weight) in examined Claris samples from EL-Bagoria canal and EL-Menofi drainage

	Diazinon				Malathion				Chlorpyrifos			
	Min	Max	Mean	St. error	Min	Max	mean	St. error	Min	Max	mean	±St. error
El-Bagoria	0	0	0.00	0.00	0	0	0.00	0.00	0	0	0.00	0.00
EL-Menofi	2.5	12	6.8	1.42	2.5	10	4.64	0.81	2	7.5	4.0	0.96




Figure 2. Mean residue levels of Organophosphorus pesticides (ppb, wet weight) in examined *Claris* samples from EL-Bagoria canal and EL-Menofi drainage

5. Discussion

There were no organophosphorus pesticides in Bahr Shabin.

The results on the table 2 and figure 1 revealed that no diazinon in Tilapia from EL-Bagoria but mean level of malathion 5.50 ± 2.0 , chlorpyrifos 5.0 ± 1.66 but from EL-Menofi drainage, mean level of diazinon 3.33 ± 0.65 , Malathion 4.98 ± 1.22 and chlorpyrifos 3.0 ± 0.71 . In *Claris* samples there was no diazinon, no Malathion, no chlorpyrifos while from EL-Menofi drainage, mean value of diazinon 6.8 ± 1.42 , 4.0 ± 0.96 .

The results of diazinon in Tilapia from EL-Menofi drainage were higher than that detected by Ibigbami et al. (2016) and (Yahia & EL-Sharkawy, 2014).

The results of Malathion in Tilapia from EL-Bagoria canal and from EL-Menofi drainage were higher than that detected by Soumis et al. (2003) and (Yahia & EL-Sharkawy, 2014).

The results of chlorpyrifos in Tilapia from EL-Menofi drainage were equal to that detected by Soumis et al. (2003) but in Tilapia from EL-Bagoria canal were higher than that detected by Soumis et al. (2003) and (Yahia & EL-Sharkawy 2014).

The results of diazinon in *Claris* from EL-Menofi drainage were higher than that detected by Ibigbami et al. (2016) and (Yahia & EL-Sharkawy, 2014).

The results of Malathion in *Claris* from EL-Menofi drainage were higher than that detected by Soumis et al. (2003) and (Yahia & EL-Sharkawy 2014).

The results of chlorpyrifos in *Claris* from EL-Menofi drainage were higher than that detected by Soumis et al. (2003) (0.3 ± 0.3) and (Yahia & EL-Sharkawy, 2014).

References

- AOAC. (1980). Official Methods of Analysis, Association Official Analytical Chemists, Washington D.C. (29 pesticides) Bej.
- Choudhary, A., Ali, A.S. and Sharique, A.A. (2014). Adverse Health Effects of Organophosphate Pesticides among Occupationally Exposed Farm Sprayers: A Case Study of Bhopal Madhya Pradesh, India. *Asian Journal of Biomedical and Pharmaceutical Sciences*, 04(35), 29–34.
- Edris, M. A.1, Fatin Said Hassanien, Fahim A. Shaltout, Azza, H.ELbaba, Nairoz, M. Adel. (2017a). Microbiological evaluation of some frozen and salted fish products in Egyptian markets. *Benha Veterinary Medical Journal*, 33(2), 317–328.
- Edris, M. A.1, Fatin Said Hassanien, Fahim A. Shaltout, Azza, H.ELbaba, Nairoz, M. Adel. (2017 b). Microbiological evaluation of some frozen and salted fish products in Egyptian markets. *Benha Veterinary Medical Journal*, 33(2), 317–328.
- FAO/WHO. (1987). Food standard programme, Codex Alimentarius Commission Room.
- Goldman, L.R., Smith, D. F, Neutra, R. R., Sounders, L. and Kiger, K. W. (1990). Pesticide food poisoning from contaminated water meloni In California, 11985, *Arch. Environ. Health*, 45,229.

- Hassan, M.A and Shaltout, F.A. (2004). Comparative Study on Storage Stability of Beef, Chicken meat, and Fish at Chilling Temperature. *Alex. J. Vet. Science*, 20(21), 21–30.
- Hassan, M. A, Shaltout, F. A., Maarouf, A. A. and El-Shafey W.S. (2014). Psychrotrophic bacteria in frozen fish with special reference to pseudomonas species. *Benha Vet. Med.J.* 27 (1), 783.
- Hassan M., Shaltout, F.A., Nabila E. El-sheikh3, Naglaa M. Sakr. (2019). Assessment of histamine residues in smoked and salted fish. Corresponding author: Naglaa Mahmoud Sakr. Animal Health Research Institute, Tanta, *Benha Veterinary Medical Journal*, 37(2), 50–52.
- Ibigbami, O.A., Aiyesanmi, A.F., Adeyeye, E.I., Adebayo, A.O. and Aladesanwa, R.D. (2016). Concentration and Potential Health Risks Associated with Organophosphorus Pesticides Residues in Fish from Three Rivers in Ekiti State, South-Western Nigeria. *African Journal of Basic & Applied Sciences*, 8(6), 324–331.
- Johnson, W. and Finley, M. T. (1980). *Handbook of acute toxicity of chemicals to fish and aquatic invertebrates*. U.S. Fish and Wildlife Service, Publication, 137. Washington, D.C., USA.
- Neuman, G. B. (1988). The occurrence and variation of Organochlorine pesticide residues detected in Australian livestock at slaughter, *Acta Veterinaria Scandinavica*, 84, 299.
- Pesticide Analytical Manual vol. I Methods which detect Multiple Residues. (1978). U.S. Department of health and Human Services. Food and Drug Administration.
- Shaltout, F. A. (2003). *Yersinia Enterocolitica* in some meat products and fish marketed at Benha city. The Third international conference Mansoura 29–30 April.
- Shaltout, F.A. and Hashim, M.F. (2002). Histamine in salted, Smoked and Canned Fish products. *Benha Vet. Med.J.* 13(1), 1–11.
- Shaltout, F.A., Hashim, M.F. and Elnahas, S. (2015). Levels of some heavy metals in fish (tilapia nilotica and Clarias lazera) at Menufia Governorate. *Benha Vet. Med.J.* 29(1), 56–64.
- Soumis, N., Lucotte, M., Sampaio, D., Al-Meida, D., C., Giroux, D, Morais, S. and Pichet, P. (2003). Presence of Organophosphate Insecticides in fish of the Amazon River. *Acta Amazonica*, 33(2), 325–338.
- Yahia, D., and Elsharkawy, E.E. (2014). Multi pesticide and PCB residues in Nile tilapia and catfish in Assiut City, Egypt. *Science of the Total Environment*, 466, 306–314.